

At page 5, please delete the three paragraphs at lines 3-34 and substitute therefor the following three paragraphs:

-- Soluble forms of RANKL are also within the scope of the invention. The nucleotide and predicted amino acid sequence of the RANKL is shown in SEQ ID Nos:11 and 13 (murine and human, respectively). Computer analysis indicated that the RANKL is a Type 2 transmembrane protein; murine RANKL contains a predicted 48 amino acid intracellular domain, 21 amino acid transmembrane domain and 247 amino acid extracellular domain, and human RANKL contains a predicted 47 amino acid intracellular domain, 21 amino acid transmembrane domain and 249 amino acid extracellular domain.

Soluble RANKL comprises a signal peptide and the extracellular domain or a fragment thereof. An exemplary signal peptide is that shown in SEQ ID NO:9; other signal (or leader) peptides are well-known in the art, and include that of murine Interleukin-7 or human growth hormone. RANKL is similar to other members of the TNF family in having a region of amino acids between the transmembrane domain and the receptor binding region that does not appear to be required for biological activity; this is referred to as a 'spacer' region. Amino acid sequence alignment indicates that the receptor binding region is from about amino acid 162 of human RANKL to about amino acid 317 (corresponding to amino acid 139 through 294 of murine RANKL, SEQ ID NO:11), beginning with an Ala residue that is conserved among many members of the family (amino acid 162 of SEQ ID NO:13).

B3  
Moreover, fragments of the extracellular domain will also provide soluble forms of RANKL. Those skilled in the art will recognize that the actual receptor binding region may be different than that predicted by computer analysis. Thus, the N-terminal amino acid of a soluble RANKL is expected to be within about five amino acids on either side of the conserved Ala residue. Alternatively, all or a portion of the spacer region may be included at the N-terminus of a soluble RANKL, as may be all or a portion of the transmembrane and/or intracellular domains, provided that the resulting soluble RANKL is not membrane-associated. Accordingly, a soluble RANKL will have an N-terminal amino acid selected from the group consisting of amino acids 1 through 162 of SEQ ID NO:13 (1 through 139 of SEQ ID NO:11). Preferably, the amino terminal amino acid is between amino acids 69 and 162 of SEQ ID NO:13 (human RANKL; amino acids 48 and 139 of SEQ ID NO:11). Similarly, the carboxy terminal amino acid can be between amino acid 313 and 317 of SEQ ID NO:13 (human RANKL; corresponding to amino acids 290 through 294 of SEQ ID NO:11). Those skilled in the art can prepare these and additional soluble forms through routine experimentation.--

At page 30, please delete in its entirety the paragraph beginning at line 36, and substitute therefor the following paragraph:

B4  
-- Addition of RANKL to DC cultures significantly increased the degree of DC aggregation and cluster formation above control cultures, similar to the effects seen with CD40L. Sorted human CD1a<sup>+</sup> DC were cultured in a cytokine cocktail (GM-CSF, IL-4, TNF- $\alpha$  and FL), in cocktail plus CD40L (1 $\mu$ g/ml), in cocktail plus RANKL (1 $\mu$ g/ml), or in cocktail plus heat inactivated ( $\Delta$ H) RANKL (1 $\mu$ g/ml) in 24-well flat bottomed culture plates in 1 ml culture media for 48-72 hours and then photographed using an inversion microscope. An increase in DC aggregation and cluster formation above control cultures was not evident when heat inactivated RANKL was used, indicating that this effect was dependent on biologically active protein. However, initial phenotypic analysis of adhesion molecule expression indicated that RANKL-induced clustering was not due to increased levels of CD2, CD11a, CD54 or CD58.--

At page 31, please delete in its entirety the paragraph beginning at line 11, and substitute therefor the following paragraph:

B5  
-- The addition of RANKL to CD1a<sup>+</sup> DC enhanced their allo-stimulatory capacity in a mixed lymphocyte reaction (MLR) by at least 3- to 10-fold, comparable to CD40L-cultured DC (Figure 2). Allogeneic T cells (1x10<sup>5</sup>) were incubated with varying numbers of irradiated (2000 rad) DC cultured as indicated above in 96-well round bottomed culture plates in 0.2 ml culture medium for four days. The cultures were pulsed with 0.5 mCi [<sup>3</sup>H]-thymidine for eight hours and the cells harvested onto glass fiber sheets for counting on a gas phase  $\beta$  counter. The background counts for either T cells or DC cultured alone were <100 cpm. Values represent the mean  $\pm$  SD of triplicate cultures. Heat inactivated RANKL had no effect. DC allo-stimulatory activity was not further enhanced when RANKL and CD40L were used in combination, possibly due to DC functional capacity having reached a maximal level with either cytokine alone. Neither RANKL nor CD40L enhanced the *in vitro* growth of DC over the three day culture period. Unlike CD40L, RANKL did not significantly increase the levels of HLA-DR expression nor the expression of CD80 or CD86.--

At page 35, please delete in its entirety the paragraph beginning at line 34, and substitute therefor the following paragraph:

B6  
Comparison of the nucleotide sequence of murine and human RANK indicated that there were several conserved regions that could be important for TRAF binding.